

Research Article

Screening of *Ocimum Gratissimum* Germplasms in Natural Field Condition against Leaf Spot Disease Caused by *Alternaria Alternata*

Digvijay Singh¹, Ravi Pratap Singh*², Ashish Kumar Singh³, Subodh Kumar Pandey¹, Hanuman Ram⁴, Amit U. Paschapur³, Manoj Parihar⁵, Rajendra Prasad⁵, and Ashutosh K. Rai⁶

¹Department of plant pathology, NDUAT, Kumarganj, Ayodhya (U.P.)

²Department of Horticulture, NDUAT, Kumarganj, Ayodhya (U.P.)

³Crop Protection Section, ICAR-VPKAS, Almora, Uttarakhand-110012, India

⁴Crop Improvement Division, ICAR-VPKAS, Almora, Uttarakhand-110012, India

⁵Crop Production Division, ICAR-VPKAS, Almora, Uttarakhand-110012, India

⁶Department of Soil Science, SVBPUAT, Meerut (U.P.)

Abstract

A field experiment was conducted during *kharif* season (2016-2017) on Tulsi (*Ocimum gratissimum*) crop for screening of resistance in natural field condition against leaf spot disease caused by *Alternaria alternata* at Main Experimental Station, Department of Medicinal and Aromatic Plants, NDUAT, Kumarganj, Faizabad. The observed parameters of oil content per cent, number of leaves per plant, leaf weight per plant (kg), leaf yield (q/ha) and disease severity per cent were recorded in 90 days old plants. The results obtained revealed that germplasm NOB-7 showed maximum oil content (2.14%) while the maximum leaves yield recorded in NOB-1 (14.81 q/ha) with disease severity (15%).

Keywords: Germplasm, oil content, leaf yield, leaf weight, disease severity, *Ocimum gratissimum*

*Correspondence

Author: Ravi Pratap Singh
Email: rp93123@gmail.com

Introduction

The Basil is native of Asia and Africa and grows wild as a perennial on some pacific islands and was brought from India to Europe through the Middle East in sixteenth century, subsequently to America in the seventeenth century.

Three types of Tulsi are encountered with in cultivation, the green leafed (Sri or Rama Tulsi) is the most common, the second type (Krishna Tulsi) bears dark green-to-purple leaves, a third type is a forest variety Vana Tulsi (*Ocimum gratissimum*) that often grows wild. *Ocimum gratissimum* is a herbaceous plant of the *Lamiaceae* family. Tulsi meaning 'the incomparable one' is an important medicinal plant which is in demand. The medicinal properties of Tulsi were known since antiquity. It is used for the treatment of problems related to heart, blood, intestine and snake bite. Eugenol the important chemical constituent of Tulsi is useful for the synthesis of vanillin.

Ocimum gratissimum leaf extract is commonly used in traditional medical practice for the treatments of mental illness, epilepsy, high fever, diarrhoea, pneumonia, cough, and conjunctivitis [1]. It has been estimated that over 50% of medicines have their origins in this natural product [2].

Material and Methods

The experiment was conducted at experimental farm of Medicinal and Aromatic Plants of Narendra Deva University of Agriculture and Technology, Kumarganj, Faizabad. Randomized Block Design (RBD) was adopted with three replications. The percent disease intensity (PDI) was recorded during August to October 2017. Twenty-four germplasm obtained from the Department of Medicinal and Aromatic Plants bearing the IC number from IC- NOB-1 to IC-NOB-24 shown in (**Table 1**) were sown in July, 2016 at Main Experimental Station, Department of Medicinal and Aromatic Plant in Randomized Block Design. The crop suffers from leaf spot disease during different stages of crops but among all the leaf spot disease amounts heavy loss in leaf yield which ultimately effects the oil yield.

The twenty-four germplasms of *Ocimum gratissimum* were screened, disease severity will be recorded using 0-9 scale. Each germplasm was planted in well prepared field at row-to-row distance 60 cm and plant to plant distance 45 cm. Details regarding the experiment are described in Table 1.

Results and discussion

The present investigation has been carried during on 2016-17 "Screening of *Ocimum gratissimum* germplasms in natural field condition against leaf spot disease caused by *Alternaria alternata*".

Table 1 List of germplasms of *Ocimum gratissimum*

S. No.	Germplasms	S.No.	Germplasms
1	NOB-1	13	NOB-13
2	NOB-2	14	NOB-14
3	NOB-3	15	NOB-15
4	NOB-4	16	NOB-16
5	NOB-5	17	NOB-17
6	NOB-6	18	NOB-18
7	NOB-7	19	NOB-19
8	NOB-8	20	NOB-20
9	NOB-9	21	NOB-21
10	NOB-10	22	NOB-22
11	NOB-11	23	NOB-23
12	NOB-12	24	NOB-24

Source: Germplasms were procured from Department of Medicinal and Aromatic plants

The initial symptoms appeared at middle leaves as small light brown, circular to sub circular spots but in some spots coalesce to form necrotic spots and cover large area. Similar symptoms were also have been reported on Tulsi crop earlier [3, 4] that the disease affects 10% of 60-day old plants and 40% of 5-month-old plants. The brown-black lesions often surrounded by a yellow halo developed from the margins and tips of the upper side of older leaves, leading to the progressive defoliation of plants, followed by plant death (**Figure 1A, B**). In later stages leaves may turn brown and die. Although *Alternaria alternata* reported on *Ocimum* sp. from Kenya [5], *Alternaria* sp. on *Ocimum bacilicum* from California [6] and Florida [7] and *Alternaria tenuissima* from Pakistan [8] and from India, *Alternaria alternata* has been reported on leaves of *Ocimum sanctum* from Poona [9], on seeds of *Ocimum* sp. from Solan, H.P. [10] and leaves of *Ocimum basilicum* from Nagarjun University, A.P. [10].




Figure 1 (A) Healthy plant of *Ocimum gratissimum*, (B) Spots of *Alternaria alternata* on *Ocimum gratissimum*, (C) Fungus of *Alternaria alternata* shown in the plate, (D) Conidia of *Alternaria alternata* on *Ocimum gratissimum*

The fungus *Alternaria alternata* was isolated from the diseased leaves of *Ocimum gratissimum*, which has some healthy portion on PDI slants. The growth of fungus was observed after five days of incubation at $25 \pm 2^\circ\text{C}$. The fungal colony was olivaceous black with dark olive-green margins, and abundant branched septate, golden-brown mycelium (Figure 1C, D). The conidiophores were branched, straight, golden-brown and smooth walled. The conidia were obpyriform, muriform produced in long branched chains, with a short pale beak [4] have observed that a fungus, consistently isolated on PDA from symptomatic leaves, formed conidia singly or in short chains (2-8 elements), dark brown, with 3-7 transverse and 0-4 longitudinal septa, $23.7-73.4 \times 8.8-15.1 \mu\text{m}$ in size, and with a conical or cylindrical beak $3.5-19.4 \mu\text{m}$ long. The pathogen was identified as *Alternaria* sp. based on morphology.

Table 2 Effect of meteorological data on development of disease on *Ocimum gratissimum* (Data recorded by Deptt. Of Agromterology, NDUAT, Kumarganj, Ayodhya)

Date of sowing	Months in which data was recorded	Standard week	Rainfall (mm)	Temperature $^\circ\text{C}$		R.H. (%)	Diseases severity (%)
				Min.	Max.		
27 July 2017	August 2017	31	20.6	26.9	32.6	85.7	35.12
		33	1.0	26.8	31.9	84.4	37.44
		34	38.6	26.3	32.9	84.7	42.53
	September 2017	35	100.6	25.6	33.0	81.3	48.23
		36	49.4	26.0	34.0	80.1	44.82
		38	88.2	25.8	32.7	84.8	46.75
	October 2017	40	0.0	24.8	33.8	79.9	67.55
		42	0.0	21.4	33.7	71.7	63.78
		43	0.0	16.7	32.7	67.5	66.00

Table 3 Correlation co-efficient of disease severity in relation to meteorological data ((Data recorded by Deptt. Of Agromterology, NDUAT, Kumarganj, Ayodhya))

Months	Rain fall (mm)	Temperature $^\circ\text{C}$		Relative Humidity (%)	Disease severity (%)
		Min.	Max.		
August 2017	0.653	-0.988	0.487	-0.574	38.36
September 2017	0.488	-0.997	-0.783	0.318	46.60
October 2017	0.0	0.323	-0.199	0.569	65.77

Table 4 Screening of *Ocimum gratissimum* germplasms in natural field condition against leaf spot disease caused by *Alternaria alternata* (Main Experiment Station Horticulture NDUAT, Kumarganj, Ayodhya)

S. No.	Entries	Disease severity (%)	No. of leaves /plant	Green leaf weight /plant (Kg)	Green leaf yield (Q/ha)	Oil content (%)
1.	NOB-1	15	2897	0.144	14.81	0.44
2.	NOB-2	13	3245	0.142	8.51	1.49
3.	NOB-3	62	1380	0.032	3.33	0.58
4.	NOB-4	44	2176	0.053	5.46	0.62
5.	NOB-5	66	2348	0.028	2.96	0.86
6.	NOB-6	37	2762	0.064	6.66	1.53
7.	NOB-7	40	2480	0.103	10.67	2.14
8.	NOB-8	20	1541	0.062	6.38	0.91
9.	NOB-9	58	2010	0.041	4.25	1.05
10.	NOB-10	30	1835	0.082	8.51	0.63
11.	NOB-11	23	1752	0.091	9.44	1.86
12.	NOB-12	35	1642	0.070	7.24	0.54
13.	NOB-13	40	1376	0.062	6.38	0.50
14.	NOB-14	45	1862	0.051	5.27	1.20
15.	NOB-15	50	1754	0.045	4.62	1.02
16.	NOB-16	55	1840	0.040	4.16	1.54
17.	NOB-17	40	2044	0.062	6.38	1.02
18.	NOB-18	18	1665	0.104	10.74	0.66
19.	NOB-19	32	1488	0.074	7.68	1.08
20.	NOB-20	38	1932	0.090	9.25	1.75
21.	NOB-21	36	2110	0.068	7.03	1.55
22.	NOB-22	60	1738	0.037	3.88	1.52

23.	NOB-23	42	2052	0.055	5.74	1.33
24.	NOB-24	26	2246	0.063	6.48	0.56

In the month of August, the disease severity was in negative correlation (-0.988) with minimum temperature while the maximum temperature was significantly positive (0.487) in relation to disease severity followed by month of September where disease severity was in negative correlation (-0.997) with the minimum temperature and maximum temperature was also negatively correlated (-0.783) to disease severity while in the month of October the disease severity was positively correlated (0.323) with minimum temperature and negative correlated (-0.199) with maximum temperature, whereas the disease severity was positive correlated with rain fall, relative humidity (RH) in rest of the months, while September and October month were significantly positive towards rain fall and relative humidity (RH).

Conclusion

The minimum percent disease intensity was recorded in month of August followed by September. The maximum percent disease intensity was noted in month of October. In the month of August, the disease intensity mean has negative correlation with minimum temperature while, the maximum temperature was significantly positive. Similarly, in month of September the disease intensity was negative correlated with minimum temperature as well as maximum temperature. In the October month disease intensity mean have significantly positive correlation with minimum and negative with maximum temperatures. On the final note keeping in view the importance of experimental crop *Ocimum gratissimum* having high medicinal value specially in Ayurveda as the herbs was known for its antiquity.

References

- [1] Eboh, DEO., Ekundina, V.O. (2013). Histological effects of chronic administration of *Ocimum gratissimum* leave extract on selected organs of adult wistar rats. *Nigerian J Sci Envir*, 12(2): 70-75.
- [2] Mc Corkle, C. M. (1995). Back to the future: Lessons from ethno veterinary research, development and extension for studying and applying local knowledge *AgricFood Hum Values Soc*, 22: 52-80.
- [3] Kumar, S., Singh, R., Chaurasia, B.M. and Kamal. (2016). First report of *Ocimum gratissimum* as a new host for pathogenic fungus *Alternaria alternata* causing Leaf spot disease in Uttar Pradesh, India. *J Pl Pathol Microbiol*, 7:361.
- [4] Garibaldi, A., Gilardi, G., Bertoldo, C. and Gullino, M.L. (2011). First report of a leaf spot of sweet Basil (*Ocimum basilicum*) caused by *Alternaria alternata* in Italy. *J Pl Pathol*, 93(4):63-89.
- [5] Caretta, G. Piontelli, E. Picco, A.M. and Del Frate, G. (1999). Some filamentous fungi on grassland vegetation from Kenya. *Mycopathologia*, 145:155-169.
- [6] French, A.M. (1989). *California Plant Disease Host Index*. Calif. Dept. Food Agric., Sacramento: 394.
- [7] Alfieri Jr, S.A., Langdon, K.R., Wehlburg, C. and Kimbrough, J.W. (1984). *Index of Plant Diseases in Florida (Revised)*. Florida Dept Agric and Consumer Serv, Div Pl Ind Bull, 11:1-389.
- [8] Ahmad, S., Iqbal, S.H. and Khalid, A.N. (1997). *Fungi of Pakistan*. Sultan Ahmad Mycolog Society Pak:248.
- [9] Narendra, D.V. and Rao, V.G. (1975). A new *Alternaria* blight of *Ocimum sanctum* from India. *Res J Mahatma Phule Agri Uni*, 6: 159-160.
- [10] Sharma, A.D. (1977). Studies on seed mycoflora of some medicinal plants. *Ind J Mycol Pl Pathol*, 7:171-172.
- [11] Vijayalaxami, M. and Rao, A.S. (1989). A new host record for *Alternaria alternata* (Fr) Keisller and its toxigenic potential. *Current Science*, 58:19.

© 2021, by the Authors. The articles published from this journal are distributed to the public under “**Creative Commons Attribution License**” (<http://creativecommons.org/licenses/by/3.0/>). Therefore, upon proper citation of the original work, all the articles can be used without any restriction or can be distributed in any medium in any form.

Publication History

Received	10.04.2020
Revised	28.05.2021
Accepted	02.06.2021
Online	30.06.2021